

A REMARK ON THE GEOMETRIC JACQUET FUNCTOR

NORIYUKI ABE AND YOICHI MIEDA

ABSTRACT. We give an action of N on the geometric Jacquet functor defined by Emerton-Nadler-Vilonen.

Let $G_{\mathbb{R}}$ be a reductive linear algebraic group over $\mathbb{R}$, $G_{\mathbb{R}} = K_{\mathbb{R}}A_{\mathbb{R}}N_{\mathbb{R}}$ an Iwasawa decomposition, and $M_{\mathbb{R}}$ the centralizer of $A_{\mathbb{R}}$ in $K_{\mathbb{R}}$. Then $P_{\mathbb{R}} = M_{\mathbb{R}}A_{\mathbb{R}}N_{\mathbb{R}}$ is a Langlands decomposition of a minimal parabolic subgroup. We use lower-case fraktur letters to denote the corresponding Lie algebras and omit the subscript “ $\mathbb{R}$ ” to denote complexifications. Fix a Cartan involution θ such that $K = \{g \in G \mid \theta(g) = g\}$. For a $(\mathfrak{g}, K)$ -module V , the Jacquet module $J(V)$ of V is defined by the space of $\mathfrak{n}$ -finite vectors in $\varprojlim_{k \rightarrow \infty} V/\theta(\mathfrak{n})^k V$ [Cas80].

For simplicity, assume that V has the same infinitesimal character as the trivial representation. Denote the category of Harish-Chandra modules with the same infinitesimal characters as the trivial representation by HC_{ρ} . Let X be the flag variety of G , $\mathrm{Perv}_K(X)$ the category of K -equivariant perverse sheaves on X . By the Beilinson-Bernstein correspondence and the Riemann-Hilbert correspondence, we have the localization functor $\Delta: \mathrm{HC}_{\rho} \rightarrow \mathrm{Perv}_K(X)$. Emerton-Nadler-Vilonen gave a geometric description of $J(V)$ by the following way [ENV04]. Fix a cocharacter $\nu: \mathbb{G}_m \rightarrow A$ which is positive on the roots in $\mathfrak{n}$. Define $a: \mathbb{G}_m \times X \rightarrow X$ by $a(t, x) = \nu(t)x$. Consider the following diagram,

$$X \simeq \{0\} \times X \rightarrow \mathbb{A}^1 \times X \leftarrow \mathbb{G}_m \times X \xrightarrow{a} X.$$

Let $R\psi$ be the nearby cycle functor with respect to $\mathbb{A}^1 \times X \rightarrow \mathbb{A}^1$. Then the geometric Jacquet functor Ψ is defined by

$$\Psi(\mathcal{F}) = R\psi(a^*(\mathcal{F})).$$

Theorem 1 (Emerton-Nadler-Vilonen [ENV04, Theorem 1.1]). *We have $\Delta \circ J \simeq \Psi \circ \Delta: \mathrm{HC}_{\rho} \rightarrow \mathrm{Perv}_K(X)$.*

It is easy to see that $J(V)$ is a $(\mathfrak{g}, N)$ -module for a $(\mathfrak{g}, K)$ -module V . Hence $\Psi(\mathcal{F})$ is N -equivariant for $\mathcal{F} \in \mathrm{Perv}_K(X)$. (See also [ENV04, Remark 1.3].)

In this paper, we give the action of N on $\Psi(\mathcal{F})$ in a geometric way. Roughly speaking, this action is given by the “limit” of the action of K .

We use the following lemma.

Lemma 2. *Let $\mathcal{X}$ be a scheme of finite type over $\mathbb{A}^1$, $\mathcal{X}^0$ (resp. $\mathcal{X}_0$) the inverse image of $\mathbb{G}_m$ (resp. $\{0\}$), and $\mathcal{G}$ a smooth group scheme over $\mathbb{A}^1$. If $\mathcal{F}^0$ is a $\mathcal{G}^0$ -equivariant perverse sheaf on $\mathcal{X}^0$, then $R\psi(\mathcal{F}^0)$ is $\mathcal{G}_0$ -equivariant.*

Proof. Define $m: \mathcal{G} \times_{\mathbb{A}^1} \mathcal{X} \rightarrow \mathcal{X}$ by $m(g, x) = gx$. Then m is a smooth morphism. Let $m^0: \mathcal{G}^0 \times_{\mathbb{G}_m} \mathcal{X}^0 \rightarrow \mathcal{X}^0$ and $m_0: \mathcal{G}_0 \times \mathcal{X}_0 \rightarrow \mathcal{X}_0$ be its restrictions. Since $\mathcal{F}^0$ is $\mathcal{G}^0$ -equivariant, we have an isomorphism $(m^0)^*(\mathcal{F}^0) \simeq \mathrm{pr}_2^*(\mathcal{F}^0)$. Hence we have

$R\psi(m^0)^*(\mathcal{F}^0) \simeq R\psi\mathrm{pr}_2^*(\mathcal{F}^0)$. By the smooth base change theorem, $m_0^*R\psi(\mathcal{F}^0) \simeq \mathrm{pr}_2^*R\psi(\mathcal{F}^0)$ [SGA7 II, Exposé XIII]. Hence $R\psi(\mathcal{F}^0)$ is $\mathcal{G}_0$ -equivariant. $\square$

Set

$$\mathcal{K}^0 = \{(t, k) \in \mathbb{G}_m \times G \mid k \in \mathrm{Ad}(\nu(t)^{-1})(K)\} \subset \mathbb{A}^1 \times G.$$

Let $\mathcal{K}$ be the closure of $\mathcal{K}^0$ in $\mathbb{A}^1 \times G$. It is a closed sub-group scheme of $\mathbb{A}^1 \times G$. Then $\mathcal{K}$ is flat over $\mathbb{A}^1$. Since each fiber of $\mathcal{K} \rightarrow \mathbb{A}^1$ is a group scheme of finite type over $\mathbb{C}$, it is reducible [Oor66]. Hence it is smooth. Therefore, $\mathcal{K}$ is smooth over $\mathbb{A}^1$.

Let Σ be the restricted root system of $(\mathfrak{g}, \mathfrak{a})$, Σ^+ the positive system corresponding to $\mathfrak{n}$, and $\mathfrak{g}_\alpha$ the restricted root space for $\alpha \in \Sigma$. Then $\mathfrak{k}$ is spanned by $\mathfrak{m}$ and $\{X + \theta(X) \mid X \in \mathfrak{g}_\alpha, \alpha \in \Sigma^+\}$. Since

$$\mathrm{Ad}(\nu(t)^{-1})(X + \theta(X)) = t^{-\langle \nu, \alpha \rangle} (X + t^{2\langle \nu, \alpha \rangle} \theta(X))$$

for $X \in \mathfrak{g}_\alpha$, the Lie algebra of $\mathrm{Ad}(\nu(t)^{-1})(K)$ is spanned by

$$\mathfrak{m} \text{ and } \{X + t^{2\langle \nu, \alpha \rangle} \theta(X) \mid X \in \mathfrak{g}_\alpha, \alpha \in \Sigma^+\}.$$

Hence the neutral component of $\mathcal{K}_0$ is $M^\circ N$ where M° is the neutral component of M . Since $MK^\circ = K$ and $\mathrm{Ad}(\nu(t)^{-1})(M) = M$, we have $\mathcal{K}_0 = MM^\circ N = MN$.

Define $\tilde{a}: \mathcal{K}^0 \times_{\mathbb{G}_m} (\mathbb{G}_m \times X) \rightarrow K \times X$ (resp. $\tilde{m}: \mathcal{K}^0 \times_{\mathbb{G}_m} (\mathbb{G}_m \times X) \rightarrow \mathbb{G}_m \times X$, $m: K \times X \rightarrow X$) by $\tilde{a}((t, k), (t, x)) = (\mathrm{Ad}(\nu(t))k, \nu(t)x)$ (resp. $\tilde{m}((t, k), (t, x)) = (t, kx)$, $m(k, x) = kx$). Then we have the following commutative diagrams

$$\begin{array}{ccc} \mathcal{K}^0 \times_{\mathbb{G}_m} (\mathbb{G}_m \times X) & \xrightarrow{\tilde{m}} & \mathbb{G}_m \times X \\ \tilde{a} \downarrow & & \downarrow a \\ K \times X & \xrightarrow{m} & X, \end{array} \quad \begin{array}{ccc} \mathcal{K}^0 \times_{\mathbb{G}_m} (\mathbb{G}_m \times X) & \xrightarrow{\mathrm{pr}_2} & \mathbb{G}_m \times X \\ \tilde{a} \downarrow & & \downarrow a \\ K \times X & \xrightarrow{\mathrm{pr}_2} & X. \end{array}$$

Let $\mathcal{F} \in \mathrm{Perv}_K(X)$. Then $m^*\mathcal{F} \simeq \mathrm{pr}_2^*\mathcal{F}$. Hence we get $\tilde{a}^*m^*\mathcal{F} \simeq \tilde{a}^*\mathrm{pr}_2^*\mathcal{F}$. By the above diagrams, $\tilde{m}^*a^*\mathcal{F} \simeq \mathrm{pr}_2^*a^*\mathcal{F}$. Therefore $a^*(\mathcal{F})$ is $\mathcal{K}^0$ -equivariant. By Lemma 2, $\Psi(\mathcal{F}) = R\psi(a^*(\mathcal{F}))$ is $\mathcal{K}_0 = MN$ -equivariant.

Remark 3. Let Γ be the quasi-inverse functor of Δ . Then N acts on $\Gamma(\Psi(\mathcal{F}))$. Moreover, this becomes a $(\mathfrak{g}, N)$ -module [Kas89, 9.1.1], namely, the infinitesimal action of N coincides with the action of $\mathfrak{n} \subset \mathfrak{g}$. We also have that $J(\Gamma(\mathcal{F}))$ is a $(\mathfrak{g}, N)$ -module. Hence both N -actions have the same infinitesimal actions. Since the action of N is determined by its infinitesimal action, the N -action we defined above coincides with the N -action on $J(\Gamma(\mathcal{F}))$.

REFERENCES

- [Cas80] W. Casselman, *Jacquet modules for real reductive groups*, Proceedings of the International Congress of Mathematicians (Helsinki, 1978) (Helsinki), Acad. Sci. Fennica, 1980, pp. 557–563.
- [ENV04] M. Emerton, D. Nadler, and K. Vilonen, *A geometric Jacquet functor*, Duke Math. J. **125** (2004), no. 2, 267–278.
- [Kas89] Masaki Kashiwara, *Representation theory and D-modules on flag varieties*, Astérisque (1989), no. 173-174, 9, 55–109, Orbites unipotentes et représentations, III.
- [Oor66] F. Oort, *Algebraic group schemes in characteristic zero are reduced*, Invent. Math. **2** (1966), 79–80.
- [SGA7 II] *Groupes de monodromie en géométrie algébrique. II*, Lecture Notes in Mathematics, Vol. 340, Springer-Verlag, Berlin, 1973, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 II), Dirigé par P. Deligne et N. Katz.

(Noriyuki Abe) GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,
3-8-1 KOMABA, MEGURO, TOKYO, 153-8914 JAPAN

E-mail address: `abenori@ms.u-tokyo.ac.jp`

(Yoichi Mieda) GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, 744 MOTOOKA,
NISHI-KU, FUKUOKA, 819-0395 JAPAN

E-mail address: `mieda@math.kyushu-u.ac.jp`